

Summary

Production Name	MEK1/2 Rabbit Polyclonal Antibody
Description	Primary antibody
Host	Rabbit
Application	WB,IHC-P,ICC/IF,IP
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide
	and 50% glycerol.
Purification	Affinity Chromatography

Immunogen

Gene Name	MAP2K1/MAP2K2
	MAP2K1; MEK1; PRKMK1; Dual specificity mitogen-activated protein kinase kinase 1;
Alternative Names	MAP kinase kinase 1; MAPKK 1; MKK1; ERK activator kinase 1; MAPK/ERK kinase 1; MEK
	1; MAP2K2; MEK2; MKK2; PRKMK2; Dual specificity mitogen-activated protein k
Gene ID	5604/5605
SwissProt ID	Q02750/P36507

Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20
Molecular Weight	Calculated MW: 43,44 kDa; Observed MW: 43,44 kDa




Background

Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Binding of extracellular ligands such as growth factors, cytokines and hormones to their cell-surface receptors activates RAS and this initiates RAF1 activation. RAF1 then further activates the dual-specificity protein kinases MAP2K1/MEK1 and MAP2K2/MEK2.


Research Area

Signal Transduction

Image Data




Western blot analysis of MEK1/2 in (1) 293T lysates; (2) A549 lysates using MEK1/2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human kidney using MEK1/2 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunofluorescence analysis of MEK1/2 in Hela using MEK1/2 antibody.

Note For research use only.