

Catalog #: APRab05619



Summary

VDR (phospho Ser51) Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Rabbit Host **Application** WB.ELISA

Reactivity Human, Mouse, Rat

Performance

Conjugation Unconjugated

Phospho Antibody Modification

Isotype IgG

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name VDR

VDR; NR111; Vitamin D3 receptor; VDR; 1; 25-dihydroxyvitamin D3 receptor; Nuclear **Alternative Names**

receptor subfamily 1 group I member 1

Gene ID 7421.0

P11473.The antiserum was produced against synthesized peptide derived from human SwissProt ID

Vitamin D3 Receptor around the phosphorylation site of Ser51. AA range:16-65

Application

Dilution Ratio WB 1:500 - 1:2000. ELISA: 1:20000

Molecular Weight 38kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: APRab05619



Background

This gene encodes the nuclear hormone receptor for vitamin D3. This receptor also functions as a receptor for the secondary bile acid lithocholic acid. The receptor belongs to the family of trans-acting transcriptional regulatory factors and shows sequence similarity to the steroid and thyroid hormone receptors. Downstream targets of this nuclear hormone receptor are principally involved in mineral metabolism though the receptor regulates a variety of other metabolic pathways, such as those involved in the immune response and cancer. Mutations in this gene are associated with type II vitamin D-resistant rickets. A single nucleotide polymorphism in the initiation codon results in an alternate translation start site three codons downstream. Alternative splicing results in multiple transcript variants encoding different proteins. [provided by RefSeg, Feb 2011], caution: It is uncertain whether Met-1 or Met-4 is the initiator, disease: Defects in VDR are the cause of type IIA rickets [MIM:277440]; also known as hypocalcemic vitamin D-resistant rickets (HVDRR). HVDRR is most frequently an autosomal recessive disorder characterized by severe rickets, hypocalcemia and secondary hyperparathyroidism.,domain:Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal steroid-binding domain., function: Nuclear hormone receptor. Transcription factor that mediates the action of vitamin D3 by controlling the expression of hormone sensitive genes. Regulates transcription of hormone sensitive genes via its association with the WINAC complex, a chromatin-remodeling complex. Recruited to promoters via its interaction with the WINAC complex subunit BAZ1B/WSTF, which mediates the interaction with acetylated histones, an essential step for VDR-promoter association. Plays a central role in calcium homeostasis, online information: The Singapore human mutation and polymorphism database, polymorphism: Genetic variations in VDR may determine Mycobacterium tuberculosis susceptibility [MIM:607948], similarity: Belongs to the nuclear hormone receptor family. NR1 subfamily, similarity: Contains 1 nuclear receptor DNA-binding domain., subunit: Homodimer in the absence of bound vitamin D3. Heterodimer with RXRA after vitamin D3 binding. Interacts with SMAD3. Interacts with MED1, NCOA1, NCOA2, NCOA3 and NCOA6 coactivators, leading to a strong increase of transcription of target genes. Interacts (in a liganddependent manner) with BAZ1B/WSTF.,

Research Area


Image Data

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab05619





Western blot analysis of lysates from COLO205 cells treated with Insulin 0.01U/ml 15 ', using Vitamin D3 Receptor (Phospho-Ser51) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of COLO cells using Phospho-VDR (S51) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA) .

Note

For research use only.