

Summary

Production Name	14-3-3-pan (Acetyl Lys51/49) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Acetyl Antibody
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	YWHAB/YWHAG/YWHAQ/YWHAZ/SFN
Alternative Names	YWHAB; 14-3-3 protein beta/alpha; Protein 1054; Protein kinase C inhibitor protein 1;
	KCIP-1; YWHAG; 14-3-3 protein gamma; Protein kinase C inhibitor protein 1; KCIP-1;
	YWHAQ; 14-3-3 protein theta; 14-3-3 protein T-cell; 14-3-3 protein tau; Protein HS1;
	YWHAZ; 14-3-3 protein zeta/delta; Protein kinase C inhibitor protein 1; KCIP-1; SFN;
	HME1; 14-3-3 protein sigma; Epithelial cell marker protein 1; Stratifin
Gene ID	7529.0
SwissProt ID	P31946.Synthesized acetyl-peptide derived from human 14-3-3-pan around the
	acetylation site of K51.

Application

```
Dilution Ratio
```

WB 1:500-1:2000. ELISA: 1:10000.

Product Name: 14-3-3-pan (Acetyl Lys51/49) Rabbit Polyclonal Antibody Catalog #: APRab06168



Molecular Weight 30kD


Background

This gene encodes a protein belonging to the 14-3-3 family of proteins, members of which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals. The encoded protein has been shown to interact with RAF1 and CDC25 phosphatases, suggesting that it may play a role in linking mitogenic signaling and the cell cycle machinery. Two transcript variants, which encode the same protein, have been identified for this gene. [provided by RefSeq, Jul 2008],function:Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathway. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner. Negative regulator of osteogenesis,,PTM:Isoform Short contains a N-acetylmethionine at position 1,,PTM:The alpha, brain-specific form differs from the beta form in being phosphorylated,,similarity:Belongs to the 14-3-3 family, subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV,,subunit:Homodimer. Interacts with SSH1 and TORC2/CRTC2. Interacts with ABL1; the interaction results in cytoplasmic location of ABL1 and inhition of cABL-mediated apoptosis. Interacts with ROR2 (dimer); the interaction results in phosphorylation of YWHAB on tyrosine residues,

Research Area

Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Oocyte meiosis;Neurotrophin;

Image Data



Western blot analysis of MOUSE-SPLEEN using Acetyl-14-3-3-pan (K51/49) antibody. Antibody was diluted at 1:500. Secondary antibody was diluted at 1:20000



Note

For research use only.