

## Summary

Production Name	DC-LAMP Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ELISA
Reactivity	Human,Rat,Mouse

### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

#### Immunogen

Gene Name	LAMP3 DCLAMP TSC403	
	Lysosome-associated membrane glycoprotein 3 (LAMP-3;Lysosomal-associated	
Alternative Names	membrane protein 3;DC-lysosome-associated membrane glycoprotein;DC	
	LAMP;Protein TSC403;CD antigen CD208)	
Gene ID	27074.0	
SwissProt ID	Q9UQV4.Synthetic peptide from human protein at AA range: 261-310	

# Application

Dilution Ratio	WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000.
Molecular Weight	44kD




## Background

Dendritic cells (DCs) are the most potent antigen-presenting cells. Immature DCs efficiently capture antigens and differentiate into interdigitating dendritic cells (IDCs) in lymphoid tissues that induce primary T-cell responses (summary by de Saint-Vis et al., 1998 [PubMed 9768752]).[supplied by OMIM, Dec 2010],function:Might change the lysosome function after the transfer of peptide-MHC class II molecules to the surface of dendritic cells.,similarity:Belongs to the LAMP family.,tissue specificity:Expressed in lymphoid organs and dendritic cells. Expressed in lung. Up-regulated in carcinomas of the esophagus, colon, rectum, ureter, stomach, breast, fallopian tube, thyroid and parotid tissues.,


### **Research Area**

Lysosome;

### Image Data




Western blot analysis of mouse-lung mouse-heart 293T Hela 3T3 lysate, antibody was diluted at 500. Secondary antibody



Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200





Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200

**Note** For research use only.