

Summary

Production Name	LPAAT-γ Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

Immunogen

Gene Name	AGPAT3	
	AGPAT3; 1-acyl-sn-glycerol-3-phosphate acyltransferase gamma; 1-acylglycerol-3-	
Alternative Names	phosphate O-acyltransferase 3; 1-AGP acyltransferase 3; 1-AGPAT 3; Lysophosphatidic	
	acid acyltransferase gamma; LPAAT-gamma	
Gene ID	56894.0	
SwissProt ID	Q9NRZ7.The antiserum was produced against synthesized peptide derived from	
	human AGPAT3. AA range:121-170	

Application

Dilution Ratio	WB 1:500-2000 ELISA 2000-20000
Molecular Weight	40kD



Background


The protein encoded by this gene is an acyltransferase that converts lysophosphatidic acid into phosphatidic acid, which is the second step in the de novo phospholipid biosynthetic pathway. The encoded protein may be an integral membrane protein. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008],catalytic activity:Acyl-CoA + 1-acyl-sn-glycerol 3-phosphate = CoA + 1,2-diacyl-sn-glycerol 3-phosphate.,domain:The HXXXXD motif is essential for acyltransferase activity and may constitute the binding site for the phosphate moiety of the glycerol-3-phosphate.,function:Converts lysophosphatidic acid (LPA) into phosphatidic acid by incorporating an acyl moiety at the sn-2 position of the glycerol backbone.,pathway:Phospholipid metabolism; CDP-diacylglycerol biosynthesis; CDP-diacylglycerol from sn-glycerol 3-phosphate: step 2/3.,similarity:Belongs to the 1-acyl-sn-glycerol-3-phosphate acyltransferase family.,

Research Area

Image Data

Glycerolipid metabolism;Glycerophospholipid metabolism;Ether lipid metabolism;

Western blot analysis of lysates from K562 cells, using AGPAT3 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from COLO205 cells using AGPAT3 antibody.



Note For research use only.