

Summary

Production Name	Olfactory receptor 2D3 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IF,WB,
Reactivity	Human, Rat, Mouse
······	

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

Immunogen

Gene Name	OR2D3
Alternative Names	OR2D3; Olfactory receptor 2D3; Olfactory receptor OR11-89
Gene ID	120775.0
SwissProt ID	Q8NGH3.The antiserum was produced against synthesized peptide derived from
	human OR2D3. AA range:231-280

Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.
Molecular Weight	36kD



olfactory receptor family 2 subfamily D member 3(OR2D3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],caution:It is uncertain whether Met-1 or Met-17 is the initiator.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


Research Area

Olfactory transduction;


Image Data




Immunofluorescence analysis of A549 cells, using OR2D3 Antibody. The picture on the right is blocked with the synthesized



Western blot analysis of lysates from HeLa and HepG2 cells, using OR2D3 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from HepG2 cells using OR2D3 antibody.





Note

For research use only.