

# Summary

Production Name	Olfactory receptor 2K2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IF,WB,
Reactivity	Human,Rat,Mouse
Host Application	Rabbit IF,WB,

# Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

# Immunogen

Gene Name	OR2K2
Alternative Names	OR2K2; OR2AR1P; Olfactory receptor 2K2; HTPCRH06; Olfactory receptor OR9-17
Gene ID	26248.0
SwissProt ID	Q8NGT1.The antiserum was produced against synthesized peptide derived from
	human OR2K2. AA range:241-290

# Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.
Molecular Weight	35kD



### Background

olfactory receptor family 2 subfamily K member 2(OR2K2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], caution: It is uncertain whether Met-1 or Met-30 is the initiator., function: Odorant receptor ,, similarity: Belongs to the G-protein coupled receptor 1 family.,


# **Research Area**

Olfactory transduction;


# **Image Data**




Immunofluorescence analysis of A549 cells, using OR2K2 Antibody. The picture on the right is blocked with the synthesized



Western blot analysis of lysates from HT-29 cells, using OR2K2 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from HepG2 cells using OR2K2 antibody.



Western Blot analysis of various cells using Olfactory receptor 2K2 Polyclonal Antibody

#### Note

For research use only.