

Summary

Production Name	Olfactory receptor 52A1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IF,WB,
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

Immunogen

Gene Name	OR52A1
Alternative Names	OR52A1; Olfactory receptor 52A1; HPFH1OR; Odorant receptor HOR3'beta4; Olfactory
	receptor OR11-319
Gene ID	23538.0
SwissProt ID	Q9UKL2.The antiserum was produced against synthesized peptide derived from human
	OR52A1. AA range:20-69

Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.

Product Name: Olfactory receptor 52A1 Rabbit Polyclonal Antibody Catalog #: APRab15256



Molecular Weight 35kD


Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


Research Area

Olfactory transduction;

Image Data




Immunofluorescence analysis of MCF7 cells, using OR52A1 Antibody. The picture on the right is blocked with the




Western blot analysis of lysates from HeLa, Jurkat, and HepG2 cells, using OR52A1 Antibody. The lane on the right is blocked





Western blot analysis of the lysates from HeLa cells using OR52A1 antibody.



Western Blot analysis of various cells using Olfactory receptor 52A1 Polyclonal Antibody

Note

For research use only.