

Catalog #: APRab17189



Summary

Ribosomal Protein S6 Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC,IF,ELISA Reactivity Human, Mouse, Rat

Performance

Conjugation Unconjugated Modification Unmodified

Isotype IgG

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name RPS6

RPS6; OK/SW-cl.2; 40S ribosomal protein S6; Phosphoprotein NP33 **Alternative Names**

Gene ID 6194.0

P62753. The antiserum was produced against synthesized peptide derived from human **SwissProt ID**

S6 Ribosomal Protein. AA range:191-240

Application

WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in

Dilution Ratio

other applications.

Molecular Weight 28kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: APRab17189




Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed function: May play an important role in controlling cell growth and proliferation through the selective translation of particular classes of mRNA, PTM: Ribosomal protein S6 is the major substrate of protein kinases in eukaryote ribosomes. The phosphorylation is stimulated by growth factors, tumor promoting agents, and mitogens. It is dephosphorylated at growth arrest, similarity: Belongs to the ribosomal protein S6e family.,

Research Area

Ribosome;mTOR;Insulin Receptor;

Image Data




Immunofluorescence analysis of HeLa cells, using S6 Ribosomal Protein Antibody. The picture on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab17189






Immunohistochemistry analysis of paraffin-embedded human skeletal muscle tissue, using S6 Ribosomal Protein Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, treated with TNF-a 20ng/ml 2 ', using S6 Ribosomal Protein Antibody. The lane on the right is blocked with the synthesized peptide.




Western Blot analysis of various cells using Ribosomal Protein S6 Polyclonal Antibody diluted at 1: 2000




Catalog #: APRab17189






Western Blot analysis of HeLa cells using Ribosomal Protein S6 Polyclonal Antibody diluted at 1: 2000 customer's (cat sample)



Western Blot analysis of customer 's (cat sample) using Ribosomal Protein S6 Polyclonal Antibody diluted at 1: 2000



Rps6

The picture was kindly provided by our customer

Note

For research use only.