

## Summary

Production Name	Wee 2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse

#### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

#### Immunogen

Gene Name	WEE2
Alternative Names	WEE2; WEE1B; Wee1-like protein kinase 2; Wee1-like protein kinase 1B; Wee1B kinase
Gene ID	494551.0
SwissProt ID	P0C1S8.The antiserum was produced against synthesized peptide derived from human
	WEE2. AA range:151-200

# Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:5000.
Molecular Weight	60kD

#### Background

### Product Name: Wee 2 Rabbit Polyclonal Antibody Catalog #: APRab19894




catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Phosphorylates and inhibits CDC2. May act as a negative regulator of entry into mitosis (G2 to M transition).,PTM:Phosphorylated .,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. WEE1 subfamily.,similarity:Contains 1 protein kinase domain.,tissue specificity:Expressed in testis.,catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Phosphorylates and inhibits CDC2. May act as a negative regulator of entry into mitosis (G2 to M transition).,PTM:Phosphorylated .,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. WEE1 subfamily.,similarity:Contains 1 protein kinase domain.,tissue specificity:Expressed in testis.,


#### **Research Area**

Cell\_Cycle\_G1S;Cell\_Cycle\_G2M\_DNA;

### Image Data



Western blot analysis of lysates from HUVEC cells, using WEE2 Antibody. The lane on the right is blocked with the



#### Western blot analysis of the lysates from K562 cells using WEE2 antibody.

#### **Note** For research use only.

