


Summary

Name	TGF- β 1/TGF-beta 1/TGFB1/Transforming Growth Factor β -1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.01 EU/ μ g as determined by LAL test.
Construction	Recombinant Human Transforming Growth Factor Beta 1 is produced by our Mammalian expression system and the target gene encoding Ala279-Ser390 is expressed.
Accession #	P01137
Host	CHO Stable Cells
Species	Human
Predicted Molecular Mass	12.8 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 50mM Glycine-HCl, 150mM NaCl, pH 2.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Alternative Names Transforming Growth Factor Beta-1; TGF-Beta-1; Latency-Associated Peptide; LAP;

Product Name: Recombinant Human TGF-beta 1
Catalog #: PCH1621



Background

TGFB1; TGFB

Transforming Growth Factor β -1 (TGF β -1) is a secreted protein which belongs to the TGF- β family. TGF β -1 is abundantly expressed in bone, articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGF β -1 performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGF β -1 peptide. TGF β -1 may also form heterodimers with other TGF β family members. It has been found that TGF β -1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.

Note

For Research Use Only , Not for Diagnostic Use.