


Summary

Name	TNF alpha/TNFSF2/TNF α
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Rabbit Tumor Necrosis Factor Alpha is produced by our E.coli expression system and the target gene encoding Val78-Leu235 is expressed.
Accession #	P04924
Host	E.coli
Species	Rabbit
Predicted Molecular Mass	17.59 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 250mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Rabbit TNF alpha
Catalog #: PEV1680



Alternative Names

Tumor Necrosis Factor; Cachectin; TNF-Alpha; Tumor Necrosis Factor Ligand Superfamily Member 2; TNF-a; TNF; TNFA; TNFSF2

Background

Tumor necrosis factor alpha (TNF α) is the prototypic ligand of the TNF superfamily. TNF α forms a homotrimer and functions by activating two types of receptors TNF-R1 (TNF receptor type 1,p55R) and TNF-R2 (TNF receptor type 2,p75R). TNF α is a pleiotropic cytokine that is capable to promote inflammation, to induce apoptotic cell death, and to inhibit tumorigenesis and viral replication. TNF α is a potent lymphoid factor that exerts cytotoxic effects on a wide range of tumor cells and certain other target cells.

Note

For Research Use Only , Not for Diagnostic Use.