


Summary

Name	NRN1L/Cpg15-2/Neuritin 1-like Protein
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Neuritin-like Protein is produced by our Mammalian expression system and the target gene encoding Ala36-Ala139 is expressed with a 6His tag at the C-terminus.
Accession #	Q496H8
Host	Human Cells
Species	Human
Predicted Molecular Mass	12.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human NRN1L (C-6His)
Catalog #: PHH1242



Alternative Names

Neuritin-like protein; NRN1L; UNQ2446/PRO5725

Background

Neuritin-like protein belongs to the neuritin family. Neuritin is a GPI-anchored protein that promotes neurite outgrowth and branching of neuritic processes in primary hippocampal and cortical cells. Neuritin expression also enhances the development of motor neuron axon arbors by promoting neuromuscular synaptogenesis and by stimulating the addition of new axon branches. Neuritin is induced by neuronal activity and by the neurotrophins, BDNF and NT3. NRN1L contains a consensus cleavage signal found in glycosylphosphatidylinositol (GPI)-anchored proteins.

Note

For Research Use Only , Not for Diagnostic Use.