
Catalog #: PHH1307



Summary

Name	Periostin(POSTN)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Osteoblast Specific Factor 2 is produced by our Mammalian expression system and the target gene encoding Asn22-Gln779 is expressed with a 6His tag at the C-terminus.
Accession #	Q15063-2
Host	Human Cells
Species	Human
Predicted Molecular Mass	85.6 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Alternative Names

Osteoblast-specific factor 2; OSF-2; POSTN; Periostin

Background Osteoblast-Specific Factor OSF-2 (POSTN, Periostin, OSF-2) is a secreted, homodimeric protein that belongs to the periostin family of the FAS1 superfamily of molecules. Periostin is a disulfide linked 90kDa bone adhesion protein secreted by osteoblasts and osteoblast-like cell lines and the protein is an attachment agent for osteoblasts. It is a TGF-beta inducible molecule that serves as both an adhesion molecule and tumor suppressor. It is synthesized by smooth muscle cells, fibroblasts and osteoblasts, as well as in the periosteum and periodontal ligament. Periostin functions as a ligand for alpha-V/beta-3 and alpha-V/beta-5 integrins to support adhesion and migration of epithelial cells.

Note

For Research Use Only, Not for Diagnostic Use.