


## Summary

<b>Name</b>	PIGR/Polymeric immunoglobulin receptor
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Polymeric Immunoglobulin Receptor is produced by our Mammalian expression system and the target gene encoding Lys19-Arg638 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	P01833
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	68.88 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human PlgR (C-6His)**  
**Catalog #: PHH1327**



---

**Alternative Names**

Polymeric Immunoglobulin Receptor; PlgR; Poly-Ig Receptor; Hepatocellular Carcinoma-Associated Protein TB6; PIGR

**Background**

The human Polymeric Immunoglobulin Receptor (plgR) is a 100 kDa type I transmembrane glycoprotein. Its precursor is 764 amino acids. It contains an 18 amino acid signal sequence, a 620 amino acid extracellular region, a 23 amino acid transmembrane fragment, and a 103 amino acid cytoplasmic domain. plgR is synthesized by secretory epithelial cells with five Ig-like domains in extracellular region, and transfer to the basolateral plasma membrane. For IgA and IgM polymers, in addition to  $\alpha$ -heavy chains and light Ig chains, a short polypeptide named joining chain (J chain) is also contained and required. plgR can bind larger polymers of IgA (plgA) and pentameric IgM as a carrier that transports IgA and IgM across epithelium. The receptor-ligand complexes are endocytosed and transcytosed to the apical surface, then proteolytic cleavage of the sixth extracellular domain of plgR and generate secretory IgA (SIgA), the plgR fragment is referred to as secretory component (SC). SIgA is a important component of the mucosal immune system. SC is anti-microbial properties and protects SIgA from proteolytic degradation

**Note**

For Research Use Only , Not for Diagnostic Use.