


## Summary

<b>Name</b>	RELT/TNFRSF19L
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Receptor Expressed in Lymphoid Tissues is produced by our Mammalian expression system and the target gene encoding Ser26-Ala160 is expressed with a human IgG1 Fc tag at the C-terminus.
<b>Accession #</b>	Q969Z4
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	41.4 KDa
<b>Formulation</b>	Supplied as a 0.2 μm filtered solution of PBS, pH7.4.
<b>Shipping</b>	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	0.00.0

## SDS-PAGE image



## Background

<b>Alternative Names</b>	Tumor necrosis factor receptor superfamily member 19L; TNFRSF19L; Receptor expressed in lymphoid tissues; RELT
<b>Background</b>	Tumor necrosis factor receptor superfamily member 19L (TNFRSF19L), also known as Receptor expressed in lymphoid tissues and RELT, is a member of the TNF-receptor superfamily. TNFRSF19L is a single-pass type membrane protein and

**Product Name: Recombinant Human RELT (C-Fc)**  
**Catalog #: PHH1423**



---

contains one TNFR-Cys repeat. TNFRSF19L is highly expressed in spleen, lymph node, thymus, peripheral blood leukocytes, bone marrow and fetal liver. It has been shown TNFRSF19L activates the NF-kappaB pathway and selectively binds TNF receptor-associated factor 1 (TRAF1). TNFRSF19L is capable of stimulating T-cell proliferation in the presence of CD3 signaling, which suggests its regulatory role in immune response.

**Note**

For Research Use Only , Not for Diagnostic Use.