


Summary

Name	IL-2/Interleukin-2/T cell growth factor/TCGF (Mutant)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Interleukin-2 Superkine is produced by our Mammalian expression system and the target gene encoding Ala21-Thr153 (L100F, R101D, L105V, I106V, I112F) is expressed with a 6His tag at the C-terminus.
Accession #	P60568
Host	Human Cells
Species	Human
Predicted Molecular Mass	16.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 4mM HCl.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in 4mM HCl. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background



Product Name: Recombinant Human IL-2 Superkine (Mutant, C-6 His)
Catalog #: PHH1918

Alternative Names

Interleukin-2; IL-2; T-Cell Growth Factor; TCGF; Aldesleukin; IL2

Background

Interleukin-2(IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system , belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes. New research has shown that IL-2 mutant reduced toxicity while being more potent at stimulating anti-tumor effector immune cells.

Note

For Research Use Only , Not for Diagnostic Use.