


# Summary

Name	PVRIG
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Transmembrane Protein PVRIG is produced by our Mammalian expression system and the target gene encoding Thr41-Asp171 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q6DKI7
Host	Human Cells
Species	Human
Predicted Molecular Mass	40.6 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### **SDS-PAGE** image



## Background



# Alternative NamesC7orf15; CD112R; PVRIG; transmembrane protein PVRIG; C7orf15MGC138295;<br/>MGC104322; MGC138297; MGC2463BackgroundHuman PVRIG (poliovirus receptor related immunoglobulin domain-containing<br/>protein), also known as CD112 receptor (CD112R), is an approximately 34 kDa<br/>single transmembrane protein in the poliovirus receptor-like protein (PVR) family.<br/>The extracellular domain sequence of human and mouse PVRIG have<br/>approximately 65% similarity. PVRIG functions as a cell surface receptor for Nectin-<br/>2/CD112, a cell surface protein that is widely expressed on antigen-presenting cells<br/>and tumor cells. Disrupting the PVRIG/Nectin-2 interaction enhances human T cell<br/>response, suggesting PVRIG is a novel checkpoint for human T cells. PVRIG may act<br/>as a coinhibitory receptor that suppresses T-cell receptor-mediated signals.

### Note

For Research Use Only, Not for Diagnostic Use.