


## Summary

<b>Name</b>	IL-17C/Interleukin-17C
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/ $\mu$ g as determined by LAL test.
<b>Construction</b>	Recombinant Human Interleukin-17C is produced by our Mammalian expression system and the target gene encoding His19-Val197 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	Q9P0M4
<b>Host</b>	Human Cells
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	20.6 KDa
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of 20mM NaAc, 50mM NaCl, 6% Trehalose, 2% Mannitol, 0.05% Tween 80, pH5.0.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human IL-17C (C-6His)**  
**Catalog #: PHH2363**



---

**Alternative Names**

CX2; Cytokine CX2; IL17C; IL-17C; IL-17CMGC126884; interleukin 17C; interleukin-17C

**Background**

Interleukin 17C (IL 17C) is a 15 20 kDa glycosylated cytokine that plays an important role in mucosal immunity and chronic inflammation. IL 17C binds to IL 17 RE with high affinity and to IL 17 RA with low affinity. These two receptor chains can associate into a heterodimeric receptor for IL 17C. IL 17C expression is induced by inflammatory stimulation in colon and airway epithelial cells, keratinocytes, CD4+ T cells, macrophages, and dendritic cells. It is up regulated in various chronic inflammatory diseases including psoriasis, cystic fibrosis, and chronic obstructive pulmonary disease (COPD). IL 17 RE is reciprocally down regulated in psoriatic lesions. The interaction of IL 17C with IL 17 RE promotes mucosal immunity through the induction of anti bacterial peptides and pro inflammatory cytokines and chemokines. IL 17C action supports the integrity of the colon epithelium following infection induced damage but also contributes to psoriatic skin thickening and the progression of arthritis. IL 17C is additionally up regulated in Th17 cell dependent autoimmunity. In this setting, it exacerbates disease severity by inducing Th17 cell production of IL 17A, IL 17F, IL 22, CCR6, and CCL20.

**Note**

For Research Use Only , Not for Diagnostic Use.