


Product Name: Recombinant Human 4-1BBL (N-6His-Flag)
Catalog #: PHH2412



Summary

Name	4-1BB Ligand/4-1BBL/CD137L/TNFSF9
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human 4-1BB Ligand is produced by our Mammalian expression system and the target gene encoding Arg71-Glu254 is expressed with a 6His, Flag tag at the N-terminus.
Accession #	P41273
Host	Human cells
Species	Human
Predicted Molecular Mass	24.5 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 50mM NaCl, 5% Trehalose, 5% Mannitol, 1mM EDTA, pH8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human 4-1BBL (N-6His-Flag)
Catalog #: PHH2412

Alternative Names Tumor necrosis factor ligand superfamily member 9; 4-1BB ligand; 4-1BBL; TNFSF9

Background Tumor necrosis factor ligand superfamily member 9(4-1BBL) is single-pass type II membrane protein which is a member of the the tumor necrosis factor family. 4-1BBL is a 254 amino acids cytokine that is expressed in brain, placenta, lung, skeletal muscle and kidney. TNFSF9 has been shown to reactivate anergic T lymphocytes in addition to promoting T lymphocyte proliferation. This cytokine may have a role in activation-induced cell death (AICD) and cognate interactions between T-cells and B-cells/macrophages. It has also been shown to be required for the optimal CD8 responses in CD8 T cells, and is thought to be involved in T cell-tumor cell interaction.

Note

For Research Use Only , Not for Diagnostic Use.