


Summary

Name	R-Spondin 3/RSPO3/Rspondin-1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human R-Spondin 3 is produced by our Mammalian expression system and the target gene encoding Gln22-Val146 is expressed with a 6His tag at the C-terminus.
Accession #	Q9BXY4
Host	Human Cells
Species	Human
Predicted Molecular Mass	14.7 kDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human RSPO3 (C-6His)
Catalog #: PHH2430



Alternative Names

R-spondin-3; RSPO3; Protein with TSP type-1 repeat; Roof plate-specific spondin-3; Thrombospondin type-1 domain-containing protein 2; PWTSR; THSD2; CRISTIN1

Background

R-spondin-3 (RSPO3), also known as Protein with TSP type-1 repeat, Roof plate-specific spondin-3, Thrombospondin type-1 domain-containing protein 2, PWTSR, THSD2 and CRISTIN1, is a member of the thrombospondin type 1 repeat supergene family. RSPO3 is a secreted protein and widely expressed in many tissues. RSPO3 contains two Furin-like repeats which have been found in a variety of eukaryotic proteins involved in the mechanism of signal transduction by receptor tyrosine kinases, and one TSP type-1 domain, RSPO3 functions as an activator of the beta-catenin signaling cascade, leading to TCF-dependent gene activation. Otherwise, RSPO3 may negatively regulate the TGF-beta pathway.

Note

For Research Use Only , Not for Diagnostic Use.