


## Summary

|                                 |   |
|---------------------------------|---|
| <b>Name</b>                     | HGF/Hepatopoietin-A/Hepatocyte Growth Factor  |
| <b>Purity</b>                   | Greater than 95% as determined by reducing SDS-PAGE   |
| <b>Endotoxin level</b>          | <0.01 EU/μg as determined by LAL test.  |
| <b>Construction</b>             | Recombinant Mouse Hepatocyte Growth Factor is produced by our Mammalian expression system and the target gene encoding Gln33-/xadLeu728 is expressed with a 6His tag at the C-terminus.   |
| <b>Accession #</b>              | Q08048  |
| <b>Host</b>                     | Human Cells   |
| <b>Species</b>                  | Mouse   |
| <b>Predicted Molecular Mass</b> | 53.5&26.9 KDa   |
| <b>Formulation</b>              | Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.   |
| <b>Shipping</b>                 | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.  |
| <b>Stability&amp;Storage</b>    | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.   |
| <b>Reconstitution</b>           | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## SDS-PAGE image



## Background

**Product Name: Recombinant Mouse HGF (C-6His)**  
**Catalog #: PHM1923**



---

**Alternative Names**

Hepatopoietin-A; Scatter factor; SF; deafness; autosomal recessive 39; DFN39; F-TCF; hepatocyte growth factor

**Background**

HGF, is a pleiotropic protein in the Plasminogen subfamily of S1 peptidases. Mouse HGF is secreted as an inactive 728 amino acid (aa) single chain propeptide. It is cleaved after the fourth Kringle domain by a serine protease to form bioactive disulfide-linked HGF with a 60 kDa alpha and 30 kDa beta chain. HGF binds heparan-sulfate proteoglycans and the widely expressed receptor tyrosine kinase, HGF R/c-MET. HGF regulates epithelial morphogenesis by inducing cell scattering and branching tubulogenesis. It can also alter epithelium morphology by the induction of nectin-1 alpha ectodomain shedding, an adhesion protein component of adherens junctions.

**Note**

For Research Use Only , Not for Diagnostic Use.