Product Name: Recombinant Mouse TrkA (C-6His)

Catalog #: PHM2345



Summary

Name TrkA/Ntrk1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse High Affinity Nerve Growth Factor Receptor is produced

by our Mammalian expression system and the target gene encoding Ala34-

Pro418 is expressed with a 6His tag at the C-terminus.

Accession # Q3UFB7

Host Human Cells

Species Mouse

Predicted Molecular Mass 42.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse TrkA (C-6His)

Catalog #: PHM2345



Alternative Names

NTRK1; NTRK-1; p140-TrkA; TRK1-transforming tyrosine kinase protein; TrkA; Trk-A; TRKAOncogene TRK; TRKTRK1

Background

TrkA (Tyrosine kinase receptor A), also known as High affinity NGF receptor, is a member of the neurotrophic tyrosine kinase receptor family that has three members. It is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Higher affinity binding of NGF requires the coexpression of TrkA with the p75 NGF receptor (NGFR), a member of the tumor necrosis factor receptor superfamily. Mutations in TRKA gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. It was originally identified as an oncogene as it is commonly mutated in cancers, particularly colon and thyroid carcinomas.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838