


Summary

Name	B7-2/CD86/T-lymphocyte activation antigen CD86
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Rabbit T-lymphocyte Activation Antigen CD86 is produced by our Mammalian expression system and the target gene encoding Ala23-Arg246 is expressed with a 6His tag at the C-terminus.
Accession #	P42071
Host	Human Cells
Species	Rabbit
Predicted Molecular Mass	26.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Rabbit B7-2 (C-6His)
Catalog #: PHV1666



Alternative Names

T-lymphocyte activation antigen CD86; Activation B7-2 antigen; CD86

Background

T-lymphocyte activation antigen CD86 (B7-2) is a glycosylated protein in the B7 family. B7 family members are transmembrane cell surface molecules that play important roles in immune activation and the maintenance of immune tolerance. It is highly expressed on activated antigen presenting cells. CD86 involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation.

Note

For Research Use Only , Not for Diagnostic Use.